
To discuss with your health care provider



Re	sure to review your plan benefits to determine your co	osts for these services				
Ro	outine health guide					
An	nual Wellness and Routine Check-up	Annually: Discuss related screening with your doctor.				
	pesity Screening: et/Physical Activity/BMI Counseling	Annually.				
CO	sion and Dental Exam (These services may not be vered by your medical benefits plan. Check your an documents).	Annually: Discuss with your doctor.				
Re	ecommended diagnostic checkups and sci	reenings for at-risk patients				
Ab	dominal Aortic Aneurysm (AAA) Check	One-time screening for ages 65 to 75 who have ever smoked of have a family history of AAA.				
Do	una Minaral Danaity Caraaning and prescribed	Warnen beginning at age CE or older, and in younger warnen				





Live a healthy lifestyle

- · Get your annual wellness exam to review your overall health and keep follow-up visits with your doctor.
- Find out if you are at risk for health conditions such as diabetes, high cholesterol, and high blood pressure.
- · Get your vaccines, preventive screenings, and labs.
- · Human Papillomavirus (HPV) vaccine 3-dose series is recommended for men and women ages 19 through 26 years if not previously vaccinated prior to age 13.
- · Talk with your doctor about the medications and over-the-counter/ vitamins you are taking to reduce side effects and interactions.
- · Get a flu vaccine every year to prevent illness and related hospitalizations.
- Get a COVID-19 vaccine to prevent severe illness and related hospitalizations. Immunocompromised people should consult their physician on the need for an additional mRNA vaccine dose.

Sources: These guidelines are recommendations from the following organizations and were not developed by Truli for Health. www.ahrq.gov, www.cdc.gov

To discuss with your health care provider



Adult (age 19+) wellness schedule
Be sure to review your plan benefits to determine your costs for these services

Guidance						
Screen/Counseling: Depression, Anxiety, Obesity, Tobacco, Alcohol, Substance Abuse, and Pregnancy	Every visit, or as indicated by your doctor.					
Fall Risk/Unintentional Injury/Domestic Violence Prevention/Seat Belt Use	Discuss exercise and home safety with your doctor.					
Medication List (including over-the-counter and vitamins) for potential interactions	Every visit, or as indicated by your doctor.					
Advance Directives/Living Will	Annually.					

Immunizations* (routine recommendation	ns)						
Tetanus, Diphtheria, Pertussis (Td/Tdap)	Ages 19+: Tdap vaccine once, then a Td booster every 10 year						
Flu (Influenza), and RSV	Flu (Ages 19+) – annually during flu season; RSV – for pregnant persons at 32 – 36 weeks; RSV – (Ages 60 – 74) – single dose based on risk factors and physician recommendation, RSV – (Age 75+) – single dose recommended by CDC.						
Pneumococcal	Ages 19 – 64: if risk factors are present; Ages 65+: 1 – 2 doses (per CDC); Ages 50+:1 dose (Florida Blue Benefits.)						
Shingles (Zoster)	Ages 50+: 2 doses Shingrix.						
Haemophilus Influenzae Type b (HIB), Hepatitis A, Hepatitis B, Meningococcal	Ages 19+: if risk factors are present.						
Human Papillomavirus (HPV), Measles/Mumps/ Rubella (MMR), Varicella (Chickenpox), and Hepatitis C (HCV) Infection Screening	Physician recommendation based on past immunization or medical history.						
COVID-19	Recommended for adults ages 19 and older within the scope of the authorization/approval for the particular vaccine.						

^{*} Some immunizations are contraindicated for certain conditions, discuss with your doctor



We're here to help:

Call

Customer Service

1-855-30-TRULI (1-855-308-5692)

TTY/TDD: Call 711

Go to TruliForHealth.com

Visit a Florida Blue Center

Go to FloridaBlueCenters.com for locations or call 1-877-352-5830

Sources: These guidelines are recommendations from the following organizations and were not developed by Truli for Health. www.ahrq.gov, www.cdc.gov

To discuss with your health care provider



Routine health guide						
Wellness Exam and Autism/Development Behavioral Assessment	Newborn up to age 3: Frequent Wellness Check- ups; Age 3 – 18: Annual Wellness Check-up.					
Body Mass Index (BMI): Height and Weight	Every visit, BMI beginning at age 2.					
Blood Pressure	Annually, beginning at age 3.					
Hearing/Dental/Vision Screenings (These services may not be covered by your medical benefits plan. Check your plan documents.)	Hearing: Newborn then annually beginning at Age 4; Dental: Regularly, beginning at age 1; Vision: Annually, beginning at age 3.					
Recommended screenings for at-risk patients						
Cholesterol Screening	Annually, beginning at age 2.					
Lead test, TB, Sickle Cell, and Blood Sugar	As indicated by history and/or symptoms.					
HIV and other Sexually Transmitted Infections (STIs) Screening and Counseling	Discuss with your doctor based on behavioral risks.					
Skin Cancer Screening	Discuss with your doctor.					
Guidance						
Injury/Violence Prevention	Annually, more often if indicated by your doctor.					
Diet/Physical Activity/Emotional Well-Being Counseling	Every visit.					
Tobacco/Alcohol/Substance Abuse/Depression/Pregnancy Screening and Counseling	Every visit starting at age 11, earlier if indicated by your doctor.					



Are your children up-todate with vaccinations?

Getting the recommended sequence of vaccinations is always a good idea to protect your child from illnesses from birth to 18 years of age. Most of these vaccinations require additional doses or boosters over time. As children grow up to become teenagers, they may come in contact with different diseases. Here are vaccines that can help protect your preteen or teen from these other illnesses and infections:

Tdap vaccine

Age 11 or 12. Protects against tetanus (lock jaw), diphtheria, and acellular pertussis (whooping cough). This is a booster shot of the same vaccine given during early childhood.

Meningococcal vaccine

(MCV4) Two doses beginning at 11 or 12 years, with a booster dose at age 16. (MenB) Two doses beginning at ages 16 – 18. Protects against meningitis, sepsis (a blood infection), and other meningococcal diseases. Children with higher risk factors may need additional doses.

Human papillomavirus

(HPV) Vaccine two or three doses over six months, beginning at ages 11 – 12. (Two doses if started before 15th birthday or three if started on or after 15). Protects boys and girls against HPV, which can lead to cancers and genital warts.

Continued next page

Sources: These guidelines are recommendations from the following organizations and were not developed by Truli for Health. www.ahrq.gov, www.cdc.gov

To discuss with your health care provider

Immunizations*	Birth	1 month	2 months	4 months	6 months	12 months	15 months	18 months	24 months	2 – 6 years	11 - 12 years	13 – 14 years	15 years	16 - 18 years
Hepatitis A					2 dose series, 12 – 23		months							
Hepatitis B	1st dose	2nd	dose		3rd dose									
Diphtheria, Tetanus, Pertussis (DTaP)			1st dose	2nd dose	3rd dose	4th dose			5th dose					
Tetanus, Diphtheria, Pertussis (Tdap)											1st dose			
Haemophilus Influenzae Type b (Hib)			1st dose	2nd dose	3rd o	d or 4th dose**								
Inactivated Poliovirus			1st dose	2nd dose	3rd dose					4th dose				
Measles, Mumps, Rubella (MMR)***					1st dose				2nd dose					
Varicela						1st dose				2nd dose				
Pneumococcal			1st dose	2nd dose	3rd dose	4th dose								
Flu (Influenza)						6 montl	ns throug	ıh 8 year	s 1 or 2 d	oses; 9 y	ears and	older 1 d	ose only	
Rotavirus			1st dose	2nd dose	3rd dose**									
Meningococcal											1st dose			Booster
Human Papillomavirus (HPV)											3 dose series			
Covid 19					6 months and older per CDC guidelines for specific vaccine									



Flu Vaccine

Every year after six months of age. Protects individuals from getting the influenza virus.

Respiratory Syncytial Virus (RSV) Injection

Respiratory syncytial virus (RSV) is the leading cause of hospitalization among U.S. infants. The CDC recommends nirsevimab for all infants aged <8 months who are born during or entering their first RSV season (October through March) and for infants and children aged 8 – 19 months who are at increased risk for severe RSV disease and are entering their second RSV season.

COVID-19 Vaccine

The American Academy of Pediatrics (AAP) recommends the COVID-19 vaccination for all children and adolescents 6 months of age and older who do not have contraindications using a COVID-19 vaccine authorized for use for their age. Any authorized COVID-19 vaccine appropriate by age and health status can be used for COVID-19 vaccination in children and adolescents.

Keep your teens safe from preventable, painful, and life-threatening diseases by staying in touch with your pediatrician's office or health clinic. Be sure to verify your benefits for preventive services.

Sources: These guidelines are recommendations from the following organizations and were not developed by Truli for Health. www.ahrq.gov, www.cdc.gov

Health coverage is offered by Truli for Health, an affiliate of Florida Blue. These companies are Independent Licensees of the Blue Cross and Blue Shield Association.

^{*} These are routine immunizations based upon cdc.gov recommendations. Range of recommended ages for catch-up or certain high-risk groups is at the doctor's discretion based on the member's family history and personal risk factors.

^{**} Dosages determined by doctor with the type of brand vaccine used.

^{***} Those in an outbreak area, aged ≥12 months who previously received ≤ 2 doses should receive a dose of mumps-virus containing vaccine.